



IL-7R α ^{lo} KLRG1^{hi} cells are not always short lived effector cells



Functional CD8⁺ T-cell populations among acute phase effector cells and memory cells can be distinguished by differences in the expression of IL-7R α and KLRG1. Cells with a IL-7R α ^{lo}/KLRG1^{hi} phenotype, previously defined as short-lived effector cells in mice, persist long after the acute phase of primary CMV infection in humans. (Source Remmerswaal *et al.* 2019)

Co-expression of IL-7R α and KLRG1 in mice can be used to define CD8 effector memory phenotypes during acute viral infections. Early effectors, short-lived effectors and precursor long term memory cells can be defined as IL-7R α ^{lo}KLRG1^{lo}, IL-7R α ^{lo}KLRG1^{hi} and IL-7R α ^{hi}KLRG1^{lo}, respectively. Remmerswaal *et*

al., aimed to determine if IL-7R α /KLRG1 co-expression profiles and functional features of CD8 memory subsets defined in murine studies correspond in persistence and functions to humans memory cells during viral infection.

Researchers studied total and viral-specific CD8 T cells phenotypes in EBV and Human-CMV seropositive healthy adults. Profiling of total non-specific T cells demonstrated that IL-7R α /KLRG1 co-expression and functions defined in mice, are similar to those observed in humans. Expression of cytotoxic molecules were associated with lack of expression of IL-7R α either in the presence or absence of KLRG1. Transcriptional profiling of bulk T cells based on IL-7R α /KLRG1 co-expression patterns showed, that KLRG1 expressing cells were associated with leukocyte activation and immune responses to bacterial antigens, whereas IL-7R α expressing cells are associated with T cell differentiation. Further suggesting that KLRG1 T cell expression is associated with increased effector functions compared to IL-7R α expressing T cells. In line with this, they also showed that KLRG1⁺ CD8 T cells that expressed cytotoxic molecules were predominantly observed in the blood but low frequencies were in the lymph node.

When researchers analysed viral-specific CD8 T cells responses, they observed differential enrichment in IL-7R α /KLRG1 co-expressing cells depending on the virus. Where, Influenza- and EBV-specific cells were predominantly enriched in IL-7R α ^{hi}KLRG1^{lo} and IL-7R α ^{hi}KLRG1^{hi}, while CMV-specific cells were enriched in IL-7R α ^{lo}KLRG1^{hi} only. This phenotype enrichment was associated with reduced expression of Granzyme B and Tbet in influenza- and EBV-specific cells compared with CMV-specific T cells.

In summary, this study highlights the importance of studying antigen-specific memory T cell functional and phenotypic profiles in the context of a specific host and/or infection/disease context. Researchers showed that

differentiation models identified in one infection and/or animal model doesn't apply to all. Where in murine models IL-7R α ^{lo}KLRG1^{hi} are short lived effector cells, while in CMV infection this phenotype is also observed in long-term memory and is the predominant memory phenotype associated with CMV-latency.

Article: Remmerswaal *et al.* 2019. [Expression of IL-7R \$\alpha\$ and KLRG1 defines functionally distinct CD8+T-cell populations in humans.](#) European Journal of Immunology

Article by Cheleka AM Mpande