


# JoAnne Flynn: BCG IV vaccination induces sterilising M.tb immunity



Protection against Mtb infection after IV BCG immunization. a, Lung inflammation (total FDG activity) and number of lung granulomas over the course of infection as measured by serial PET-CT scans. Each line shows one NHP over time; 3 NHPs (2 unvaccinated (unvax) and 1 IDlow) reached a humane end point before 12 weeks. tntc, too numerous to count. b, Three-dimensional volume renderings of PET-CT scans of each NHP at the time of necropsy. PET was limited to the thoracic cavity; the standardized uptake value colour bar is shown in the top right and indicates FDG retention, a surrogate for inflammation. (Source Darrah et al., 2020)

Almost 100 years since the development of Bacille Calmette-Guerin (BCG) vaccine, it is still the only licenced vaccine against Tuberculosis (TB). BCG is predominantly given to infants, where it provides protection against disseminated TB in children but has very variable efficacy against TB in adolescents and adults.

Tuberculosis vaccinologists suggest that reduced efficacy in adults could be partly due to the route of BCG delivery. Where intradermal BCG vaccination, predominant route of delivery, does not induce sufficient long-term lung-resident memory responses. In line with this, animal TB models have demonstrated better survival post-*Mycobacterium tuberculosis* (M.tb) challenge following aerosol or intravenous BCG vaccination, compared with BCG intradermal vaccination.

In the final segment of Dr Flynn's webinar, she highlighted research (Darrah et al.,) that aimed to determine how dose and route of BCG vaccination influence the generation of systemic and tissue-resident immunity, as well as "protect" against TB. In their study, they vaccinated non-human primates (NHPs) (*Macaca mulatta*; highly susceptible rhesus macaques) with one of the following strategies:

- Normal dose intradermal vaccination: ID<sub>low</sub>
- High dose intradermal vaccination: ID<sub>high</sub>
- Aerosol vaccination (AE)
- Intravenous vaccination (IV)
- Combination of ID<sub>low</sub> and AE

Darrah et al., showed that BCG-IV resulted in superior T cell and antigen-presenting cell recruitment to the lung compared to other vaccination strategies. In addition to T cell and innate immunity, BCG-IV also induced superior vaccine-induced antibody responses. Following M.tb challenge, the BCG-IV cohort consistently had lower lung inflammation post M.tb challenge, the lowest bacterial burden with 6 out of 10 NHPs with undetectable bacterial burden and 9 out 10

protected from TB. In addition to the low M.tb burden, BCG-IV was the only vaccination strategy that did not result in an increase of Mycobacterial-specific T cell immune responses post M.tb infection. Based on these results, one of the proposed mechanisms of BCG-IV protection is due to the induction of robust systemic and lung resident T cell immunity, which sufficiently controls and eliminates M.tb burden.

It should be noted this BCG-IV vaccination model utilised very high doses of BCG ( $5 \times 10^7$  CFUs). Dr Flynn ended her webinar with unpublished data that illustrate protective BCG-IV is observed with much lower doses across a 500 fold BCG dose. Overall this data demonstrates that protection against TB infection or disease is possible

Journal Article: [Darrah et al., 2020. Prevention of tuberculosis in macaques after intravenous BCG immunization. Nature](#)

*Summary by Cheleka AM Mpande*

**This article is an adaptation of a previously published article.**